

Egyptian Herbal Monograph

Volume 1

Wild Medicinal Plants

Egyptian Drug Authority (EDA)

2025



Egyptian Herbal Monograph

Wild Medicinal Plants

Citrullus colocynthis L.

حنظل – حنضل

1. Names & Synonyms (1-3)

Citrullus colocynthis L.

Family: Cucurbitaceae.

Syns. *Cucumis colocynthis*

Colocynthis vulgaris.

Arabic: Hanzal حنظل - Handal حنضل.

English: Colocynth, Bitter apple, Bitter gourd.

2. Geographical distribution

Common in nearly all the phytogeographical regions of the country (3).

3. Parts used for medicinal purpose

Colocynth; Pulp of the peeled fruit (dried unripe, but fully grown fruit deprived of its seeds and hard outer part of pericarp) (4, 5); seeds, leaves, and roots are also used (2, 6).

4. Major chemical constituents (3)

- **Cucurbitacins and its glycosides** (7): Cucurbitacin A, B, C, D, E, I, J, K and L (8-13), 2-O- β -D-glucopyranosyl-cucurbitacin I, 2-O- β -D-glucopyranosyl-cucurbitacin L (14), colocynthein, colocynthetin (15), and Cucurbitane-type triterpenoid glycoside (saponin) (16) as colocynthosides A and B (11) are major constituents in the fruit.
- **Flavonoids:** Quercetin (leaf, stem, fruit, root), flavone-C-glucoside (Isovitexin), isoorientin and isoorientin 3'-O-methylether (fruit), C-*p*-hydroxybenzyl derivatives as 8-C-*p*-hydroxybenzylisovitexin, 6-C-*p*-hydroxybenzylvitexin, 8-C-*p*-hydroxybenzylisovitexin 4'-O-glucoside (aerial parts) (14, 17,18), kaempferol (19), catechin, myricetin (19-21), isoscoparin and isosaponarin (14).

- **Phenolic acids:** Gallic (20), *p*-hydroxybenzoic, chlorogenic, caffeic, vanillic acid, *p*-coumaric, sinapic and ferulic acids from fruit pulp (19), and 3-O-caffeoylquinic acid from the leaves (22).
- **Alkaloids:** Alkaloids from the whole fruit and pulp (13, 23- 26); choline from fruit pulp (24), 2-(nonan-8-one) -(1H)-4-quinolone and 2-(nonan-8-one) 4-methoxy-quinoline from the aerial parts (27).
- **Fatty acids:** Linoleic, oleic, palmitic, stearic, myristic, linolenic (17, 28-31) and arachidic acids from seeds (20).
- **Amino acids:** Arginine (in pulp), aspartic acid (in rind) and glutamic acid (in seeds) (32).
- **Others:** (17)
 - Protein: rich in lysine, leucine, sulfo-amino acids as methionine.
 - Vitamins: Vitamin B (Thiamine, Riboflavin and Niacin).
 - Minerals: Ca, Mg, Mn, K, P, Fe and Zn .
 - Tocopherols and Carotenes: α -Tocopherol (20), γ -tocopherol and β -carotene from fruit seed oil (33).
 - Volatile compounds (fruit pulp): 2-Methyl, 4-heptanone, 3-methyl, 2-heptanone, trimethylsilyl methanol and 1-propoxy pentane (34).

5. Medicinal uses

A. Well-established

Oral: Purgative (4-5).

B. Traditional use

Externally: Rheumatic disease (low back, knee and joints pain) (35-37).

C. colocynthis is a traditional medicinal plant for use in the specified indications exclusively based upon long-standing use.

6. Herbal preparations correlated to medicinal use

1. Oral dried fruits powder (4-5).
2. Topical Poultice (with warm cooking oils) (35-37).

7. Posology and method of administration correlated to medicinal use

Preparation 1

Indication A

Oral: 0.1 – 0.3g of dried fruits powder as purgative (4-5).

Preparation 2**Indication B**

Externally: Poultice is made of colocynth with warm cooking oils to place it topically on the joint for rheumatic pain (35-37).

Method of administration: Oral and external use.

8. Contraindications

Hypersensitivity to active substances and to other plants of the same family.

9. Special warnings and precautions for use

- Colocynth should be used under medical supervision and in proper dose as it is severely poisonous. It has a strongly irritating and painful effect on mucous membranes due to its content of cucurbitacin glycosides, out of which cucurbitacins are released in watery environments (38).
- **Diabetes:** Colocynth might lower blood sugar levels. Blood glucose levels should be regularly monitored in diabetics (39).
- **Surgery:** Colocynth might interfere with blood sugar control during and after surgery. Colocynth should be stopped at least 2 weeks before a scheduled surgery (39).
- Encouraging use of seeds (containing no saponin) instead of the whole fruit or pulp (40).

10. Interactions with other medicinal products and other forms of interaction (39)**- Digoxin**

Colocynth is a stimulant laxative which can decrease potassium levels in the body therefore can increase the risk of side effects of digoxin.

- Warfarin

Colocynth can work as a laxative. In some people colocynth can cause diarrhea which can increase the effects of warfarin and increase the risk of bleeding.

- Diuretic drugs

Colocynth is a laxative agent and can decrease potassium in the body. Diuretic drugs can also decrease potassium in the body. Accordingly, concomitant use with diuretic drugs might severely decrease potassium in the body. Some diuretic drugs that can decrease potassium include chlorothiazide, chlorthalidone, furosemide, hydrochlorothiazide, and others.

11. Fertility, pregnancy and lactation

Colocynth is not recommended in pregnancy and breast-feeding (39) <<see section 15; Relevant biological activities>>.

Long term exposure to *C. colocynthis* L. may cause adverse effects on the reproductive system and fertility (41) <<see section 15; Relevant biological activities >>.

12. Effects on ability to drive and use machines

No studies on the effect on the ability to drive and use machines have been performed.

13. Undesirable effects

- If adverse reactions occur, a doctor or a pharmacist should be consulted.
- Gastrointestinal disorders such as diarrhea, painful cramp (42), colitis (43, 44); intestinal damage (45); the dried fruit pulp is irritating to the eye and powdered colocynth pulp causes severe pain if it meets the nasal mucous membrane (46-47); leucocytosis (48-49).

14. Overdose

- Vomiting, bloody diarrhea, colic, and kidney irritation following the intake of toxic dosages (0.6 to 1 g), and then increased diuresis that progresses to anuria. Lethal dosages (starting at 2 g) lead to convulsions, paralysis and, if untreated, lead to death through circulatory collapse. The treatment of poisonings should proceed symptomatically following gastric lavage (38).
- In case of poisoning, a dilute tannic acid solution should be taken, followed by large quantities of drinks that contain eggs (albuminous drinks) (39).

15. Relevant biological activities

Analgesic, anti-inflammatory and antinociceptive

- Carrageenan-induced edema in a rat hind paw was carried out to evaluate the topical anti-inflammatory effect of the *C. colocynthis* (CC) fruit extract cream (2–8%) and the tissue levels of IL-6 and TNF- α were estimated by using a commercial ELISA kit. Also the topical antinociceptive activity of the CC cream (2–8%) was evaluated in rat formalin test. The study indicated that topical application of CC cream possesses significant anti-inflammatory and antinociceptive activities in animal models, which were probably mediated by opioid receptors and the suppression of pro-inflammatory cytokines (TNF- α and IL-6) (50).

- *C. colocynthis* aqueous extracts of the plant different parts were screened for analgesic and anti-inflammatory activities using the acetic acid writhing test in mice and the carrageenan-induced paw edema assay in rats, respectively. All extracts displayed analgesic and anti-inflammatory activities at different doses without inducing acute toxicity. The best results were obtained with immature fruits followed by seeds. The stem and root extracts were shown to possess the less significant inhibitory activity against the analgesic and anti-inflammatory models used (51). Also, the same assessment was carried out on the immature fruit and seeds organic extracts (petroleum ether, chloroform, ethyl acetate, acetone and finely methanol extract). All extracts displayed an important analgesic and anti-inflammatory activities at different doses without inducing any side effects. Experiment results provide scientific insight into the ancient practice of utilizing *CC* as analgesic and as anti-inflammatory agents (52).
- The extract of the dried pulp of the plant fruits was studied for its effects through inhibition of inflammatory cytokines secreted in obesity conditions on male mice. The fed animals received 50 mg/kg of hydroalcoholic extract by gavage for 42 days. TNF- α , IL-6 and IL-10 in serum were assayed by ELISA technique after every two weeks. The extract dramatically decreased expression of TNF- α 44.83 (** $p < 0.001$), IL-6 30.23 (** $p < 0.001$) and marginally increased IL-10 5.31 (ns- $p > 0.05$) in obese mice. This study demonstrated that, although the extract did not show anti-obesity effects, it could have an anti-inflammatory effect through down regulation of obesity-associated pro-inflammatory cytokines (53).
- The effect of ethanol extract of the roots at dose of 20 ng/ml on inflammatory cytokine expression in inflamed cells with LPS20 both cartilage cells/macrophage was examined. The results indicated that the ethanol extract of root can reduce expression levels of pro-inflammatory cytokines in inflamed cells caused by situation same as that of osteoarthritis (54).
- Sodium carboxymethyl cellulose (5%) topical gel formulations containing 3% of *Colocynth* extract, hydrolyzed extract, or acetylated extract were studied for their release through cellophane membrane and their permeability through hairless mouse skin and, for the *in vivo* anti-inflammatory activity of the different types of *colocynth* extract using the carrageenan induced paw edema model in albino rats in comparison with the commercial Voltarin Emulgel®. The acetylated extract gel showed comparatively rapid permeability through hairless mouse skin, with low release rate through cellophane membrane. The pharmacological screening revealed that the percent reduction of edema produced by *colocynth* extract was 45.39%, the hydrolyzed extract produced 54.11% inhibition and the acetylated extract produced 64.95%, while Voltarin Emulgel® produced 63.35%. This means that acetylated *colocynth* extract can be used as an effective local anti-inflammatory agent (55).

- Fruit extracts in methanol were subjected to check anti-inflammatory activity against carrageenan induced paw edema, serotonin induced edema and prostaglandin E1-induced paw edema in albino rats. Extracts showed anti-inflammatory activity against all types of edema but the most significant results were seen against prostaglandin E1-induced paw edema (56).

Toxicity

- The study was undertaken to determine the acute median lethal dose of the methanol extract of the fruit of *C. colocynthis* and to evaluate the toxic effects of this extract at a single daily oral dose (131mg/kg) in 50 Albino rats. Liver, kidney and bone marrow function test were assessed using standard techniques. The acute median lethal dose of the extract was found to be 1311.45 mg/kg. The plasma ALT, AST, urea, and creatinine levels were significantly affected, an indication that the extract is hepatonephrotoxic. The results obtained for hematological parameters reflect that methanol extract at a dose of 131 mg/kg did not affect quantitatively but disrupted qualitatively some functions of the bone marrow. The study also showed that the intake of extract of ripe *C. colocynthis* fruit presented some adverse effects on the functions of the liver, kidney, and bone marrow in rats (57).

- The toxic effects of *C. colocynthis* on male rabbits were investigated. Test animals were treated with 100 or 200 mg/kg/day of either pulp or seed extract. One month later, surviving animals were sacrificed and specimens of small intestine, kidney, and liver were prepared for morphological evaluation. No animals treated with 200 mg/kg/day of pulp extract survived. Animals treated with 100 mg/kg/day of pulp extract displayed sever lesions in the small intestine, kidney, and liver. Interestingly, animals treated with either 100 or 200 mg/kg/day of seed extract displayed only minor intestinal insult. In contrast to seeds extract, pulp extract of *C. colocynthis* can be fatal to rabbits. Therefore, seeds extract may be the preferred route for therapeutic application (45).

- The toxic effect of alcoholic extract of *C. colocynthis* on rat Liver was investigated with a single daily dose of (50, 100, 200, 400 g/kg) administered intraperitoneally. The results indicated that there is a morphological change in liver cells including karyrrhexis, chromatolysis, and granulation of the cytoplasm. Additionally, collagen and reticular fibers were evident in liver parenchyma in high doses. *C. colocynthis* can have toxic effects on liver cells which may induce hepatocyte necrosis and liver fibrosis. These effects were dose dependent (58).

- The acute toxicity studies of the methanolic extract of dried fruit pulp of *C. colocynthis* (Cucurbitaceae) in albino mice NMRI indicated the highly toxic nature of the colocynth. A very significant decrease in body weight of test animals was noted at $p < 0.05$. The LD₅₀ was calculated as 1000mg/kg body weight. Within four days of experimentation, mortality was 100%. Histopathological studies confirmed the toxic

nature of extract. Gross changes in histology of heart, liver and kidneys were noted. Section of spleen did not exhibit any abnormality (59).

- Ethanolic extract of the plant fruits was investigated for its sub-chronic toxicity on Sprague Dawley (SD) rats, to develop safe doses, 30 male rats were used with 0, 12.5 and 25 mg/kg of the extract twice per week for 8 weeks. The raw extract of the fruits at dose of 12.5 mg and 25 mg/kg induced significant increase in level of creatinine and total protein and increased non-significantly levels of glucose and blood urea while reduced level of triglycerides, total cholesterol, and high-density lipoprotein-cholesterol significantly when compared with control group. Histopathological examination revealed that kidney of rat treated with *C. colocynthis* at dose of 12.5 mg/kg show interstitial chronic inflammatory cell infiltrate while kidney of SD-rat treated with *C. colocynthis* at dose of 25 mg/kg showed interstitial fibrosis stained with Mallory trichrome stain. While liver showed lytic necrosis replaced by leukocyte infiltration at dose of 12.5 mg/kg of raw extract and rat treated with 25 mg/kg show interstitial fibrosis stained with Mallory trichrome stain. Raw extract of the fruits had adverse effect on liver and kidney while in improving lipid profile and hematology in a novel manner in SD rats at selected dosed (60).

Case Presentations

- Four patients with colocynth intoxication are presented. The main clinical feature was acute rectorrhagia preceded by mucosal diarrhea with tenesmus, which gradually progressed to bloody diarrhea and overt rectorrhagia within 3 to 4 hours. The only colonoscopic observation was mucosal erosion which was completely resolved in follow-up colonoscopy after 14 days.

The membranolytic activity of some *C. colocynthis* ingredients is responsible for the intestinal damage. Patients and herbalists should be acquainted with the proper use and side effects of the herb. Clinicians should also be aware of *C. colocynthis* as a probable cause of lower GI bleeding in patients with no other suggestive history, especially diabetics (40).

- Three examples were reported of toxic acute colitis which occurred after ingestion of *C. colocynthis* for ritual purposes. The prominent clinical feature was dysenteric diarrhea; colonoscopic changes included congestion and hyperaemia of the mucosa with abundant exudates but no ulceration or pseudopolyp formation. A causal relationship between colonic injury and the intake of colocynth was supported by the following features: (1) the pharmacology of the colocynth extract ingested; (2) the temporal relationship between colocynth intake and clinical onset (eight to 12 h); (3) the rapid recovery within three to six days, with normal endoscopy at day 14; (4) the absence of other possible causes for the observed patterns, except in one case, in which a concomitant intestinal infection with *Clostridium perfringens* Type A was discovered; (5) the specific pathological features. Colonic biopsies taken 27, 44, and 72

h after colocynth intake showed: erosions with fibrino-purulent exudate, early fibrosis of the lamina propria, hyaline thickening of the superficial epithelial basal membrane. These pathological features completely disappeared within 14 days in all three cases (61).

Pregnancy and Lactation

In Ayurvedic medicine, the fruit pulp has a reputation of causing miscarriage, when administered to pregnant women (62) and colocynth has also been used for this purpose in Europe (49, 63). Such activity could arise indirectly from congestion in the pelvic region as a manifestation of the cathartic action (62). It should be added, however, that on several occasions colocynth was ineffective as an abortive agent, even though it produced serious poisoning (63).

Fertility

- An ethanolic extract of *C. colocynthis* seeds, administered at an oral dose of 200 mg/kg for 2 days, did not inhibit copper acetate induced ovulation in rabbits to such an extent that further research seemed warranted (64). Different extracts of *C. colocynthis* were screened for anti-implantation activity by feeding female rats with each extract from day 1 to day 7 of their pregnancy. Acetone and methanolic root extracts in doses of 150 mg/kg prevented implantation in 3 and 4 of 7 test animals, respectively, whereas 200 mg/kg of an ethanolic leaf extract and 150 mg/kg of a benzene leaf extract inhibited implantation in 4 of 6 rats (65). The spermatotoxicity was observed in mice treated with an alcoholic extract of *C. colocynthis* fruit in daily oral doses of 0.1 g/kg body weight for 3 months (48). Early textbooks claim that *C. colocynthis* is excreted into breast milk and should therefore not be given to nursing women (66-68).

The short and long effects of *C. colocynthis* L. (400 mg/kg/body weight) on the reproductive system after administration to female Sprague-Dawley rats were investigated. The rats were intraperitoneally injected in dose of 400 mg/kg/body weight. First group received treatment for 4 weeks and a second group received the same dose of treatment for a period of 12 weeks. Female rats were allowed mating with males after 10 days prior to the last administration dose. Several parameters were determined including: number of pregnant rats, body and reproductive organ weight, number of implantation sites, viable fetuses and resorption sites. The results indicate that long-term exposure of female rats to *C. colocynthis* L. causes adverse effects on the reproductive system and fertility (41).

16. Additional Information

Colocynth has a drastic purgative and irritant action and has been superseded by less toxic laxatives. It is used in homoeopathic medicine (69).

C. colocynthis has showed wide range of pharmacological activities including:

- Antidiabetic, hypoglycemic and antihyperglycaemic activities (30, 70-85).
- Peripheral neuropathy (86).
- Antioxidant activity (19, 87-90).
- Hair growth effect (91, 92).
- Anti-microbial activity (13, 25, 93-98).
- Anthelmintic activity (99-103).
- Cytotoxic activity (87, 104-106).
- Hypolipidemic effect /antihyperlipidemic (89, 107,108).
- Hepatoprotective effect (109-111).

17. Date of compilation/last revision

30/8/2023.

References

1	Boulos, L. (2000). Flora of Egypt, Al Hadara Publishing, Cairo, Egypt.
2	Batanouny, K. H. (1999). Wild Medicinal Plants in Egypt. (with contribution: Aboutabl, E., Shabana, M. and Soliman, F.). Academy of Scientific Research and Technology, Egypt. The World Conservation Union (IUCN).
3	Hassan, N. M and Omer, E. A. (2018). <i>Citrullus colocynthis</i> L. In: Egyptian Encyclopedia of Wild Medicinal Plants, 6 , 18-36. Academy of Scientific Research and Technology, Cairo, Egypt.
4	Egyptian Pharmacopoeia (1984). 3 rd edition. General Organization for Government Printing. Cairo.
5	Egyptian Pharmacopoeia (2005). 4 th edition. General Organization for Government Printing. Cairo.
6	Hammouda, F. M., Ismail, S. I., Abdel-Azim, N. S. and Shams, K. A. (2005). A Guide to Medicinal Plants in North Africa (Batanouny K. H., editor). IUCN Centre for Mediterranean Cooperation. Malaga.
7	Hussain, A. I., Rathore, H. A., Sattar, M. Z. A., Chatha, S. A. S., Sarker S. D. and Gilani, A. H. (2014). <i>Citrullus colocynthis</i> (L.) Schrad (bitter apple fruit): A review of its phytochemistry, pharmacology, traditional uses and nutritional potential. <i>J. Ethnopharmacol.</i> , 155 , 54-66.
8	Adam, S. E. I., Al-Yahya, M. A. and Al-Farhan, A. H., (2001). Response of Najdi sheep to oral administration of <i>Citrullus colocynthis</i> fruits, <i>Nerium leander</i> leaves or their mixture. <i>Small Ruminants Research</i> , 40 , 239-244.
9	Chen, J. C., Chiu, M. H., Nie, R. L., Cordell, G. A. and Qiu, S. X. (2005). Cucurbitacins and cucurbitane glycosides: structures and biological activities. <i>Natural Product Reports</i> , 22 (3), 386-399.
10	Tannin-Spitz, T., Grossman, S., Dovrat, S., Gottlieb, H. E. and Bergman, M. (2007). Growth inhibitory activity of cucurbitacin glucosides isolated from <i>Citrullus colocynthis</i> on human breast cancer cells. <i>Biochemical Pharmacology</i> , 73 , 56-67.
11	Yoshikawa, M., Morikawa, T., Kobayashi, H., Nakamura, A., Matsuhira, K., Nakamura, S. and Matsuda, H. (2007). Bioactive saponins and glycosides, XXVII. Structures of new cucurbitane-type triterpene glycosides and antiallergic constituents from <i>Citrullus colocynthis</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 55 , 428-434.
12	Torkey H. M., Abou-Yousef H. M., Azeiz, A. and Farid, H. E. A. (2009). Insecticidal effect of cucurbitacin E glycoside isolated from <i>Citrullus colocynthis</i> against <i>Aphis craccivora</i> . <i>Australian Journal of Basic and Applied Sciences</i> , 3 , 4060-4066.
13	Ali, A. A., Alian, M. A. and Elmahi, H. A. (2013). Phytochemical analysis of some chemical metabolites of Colocynth plant (<i>Citrullus colocynthis</i> L.) and its activities as antimicrobial and antiplasmodial. <i>Journal of Basic and Applied Scitific Research</i> , 3 , 228-236.
14	Delazar, A., Gibbons, S., Kosari, A., Nazemiyeh, H., Modarresi, M., Nahar, L. and Sarker, S. (2006). Flavone c-glycosides and cucurbitacin glycosides from <i>Citrullus colocynthis</i> . <i>DARU J. Pharmaceut. Sci.</i> , 14 (3), 109-114.

15	Pravin, B., Tushar, D., Vijay, P. and Kishanchnad, K. (2013). Review on <i>Citrullus colocynthis</i> . <i>International Journal of Research in Pharmacy and Chemistry (IJRPC)</i> , 3 (1), 46-53.
16	Song, F., Dai, B., Zhang, H., Xie, J., Gu, C. and Zhang, J. (2015). Two new cucurbitane-type triterpenoid saponins isolated from ethyl acetate extract of <i>Citrullus colocynthis</i> fruit. <i>Journal of Asian Natural Products Research</i> , 17 (8), 1-6.
17	Gurudeeban, S., Satyavani, K. and Ramanathan, T. (2010). Bitter apple (<i>Citrullus colocynthis</i>): an overview of chemical composition and biomedical potentials. <i>Asian Journal of Plant Sciences</i> , 9 , 394-401.
18	Maatooq, G. T., El-Sharkawy, S. H., Afifi, M. S. and Rosazza, J. P. N. (1997). C-p-hydroxybenzoyl glycoflavones from <i>Citrullus colocynthis</i> . <i>Phytochemistry</i> , 44 , 187-190.
19	Hussain, A. I., Rathore, H. A., Sattar, M. Z. A., Chatha, S. A. S., Ahmad, F., Ahmad, A. and Johns, E. J. (2013). Phenolic profile and antioxidant activity of various extracts from <i>Citrullus colocynthis</i> (L.) from the Pakistani flora. <i>Industrial Crops and Products</i> , 45 , 416-422.
20	Al-Snafi, A. E. (2016). Chemical constituents and pharmacological effects of <i>Citrullus colocynthis</i> - A review. <i>IOSR Journal of Pharmacy</i> , 3 (6), 57-67.
21	Meena, M. C. and Patni, V. (2008). Isolation and identification of flavonoid "Quercetin" from <i>Citrullus colocynthis</i> (Linn.) Schrad. <i>Asian Journal of Experimental Sciences</i> , 22 , 137-142.
22	Shawkey A. M., Rabeh M. A. and Abdellatif A. O. (2014). Biofunctional molecules from <i>Citrullus colocynthis</i> : An HPLC/MS analysis in correlation to antimicrobial and anticancer activities. <i>Advances in Life Science and Technology</i> , 17 , 51-61.
23	Lahfa, F. B., Azzi, R., Mezouar, D. and Djaziri, R. (2017). Hypoglycemic effect of <i>Citrullus colocynthis</i> extracts. <i>Phytothérapie</i> , 15 , 50-56 .
24	Sayed, D. M., Balbaa, S. I. and Afifi, M. S. A. (1973). Nitrogenous bases of the different organs of <i>Citrullus colocynthis</i> . <i>Planta Medica</i> , 24 (3), 260-265.
25	Najafi, S., Sanadgol, N., Nejad, B. S., Beiragi, M. A. and Sanadgol, E. (2010). Phytochemical screening and antibacterial activity of <i>Citrullus colocynthis</i> (Linn.) Schrad against <i>Staphylococcus aureus</i> . <i>Journal of Medicinal Plants Research</i> , 4 (22), 2321-2325.
26	Mukherjee, A. and Patil, S. D. (2012). Effects of alkaloid rich extract of <i>Citrullus colocynthis</i> fruit on <i>Artemia salina</i> and human cancerous (MCF-7 and HEPG-2) cells. <i>Journal of Pharma Sci. Tech.</i> , 1 , 15-19.
27	Salama, H. M. H. (2012). Alkaloids and flavonoids from the air dried aerial parts of <i>Citrullus colocynthis</i> . <i>Journal of Medicinal Plants Research</i> , 6 (38), 5150-5155.
28	Sawaya, W. N., Daghir, N. J. and Khan, P. (1983.) Chemical characterization and edibility of the oil extracted from <i>Citrullus colocynthis</i> seeds. <i>Journal of Food Science</i> , 48 , 104-106.
29	Sadou, H., Sabo, H., Alma, M. M., Saadou, M. and Leger, C. L. (2007). Chemical content of the seeds and physico-chemical characteristic of the seed oils from <i>Citrullus colocynthis</i> , <i>Coccinia grandis</i> , <i>Cucumis metuliferus</i> and <i>Cucumis prophetarum</i> of Niger. <i>Bulletin of the Chemical Society of Ethiopia</i> , 21 , 323-330.

30	Sebbagh, N., Cruciani-Guglielmacci, C., Ouali, F., Berthault, M. F., Rouch, C., Sari, D. C. and Magnan, C. (2009). Comparative effects of <i>Citrullus colocynthis</i> , sun flower and olive oil-enriched diet in streptozotocin-induced diabetes in rats. <i>Diabetes and Metabolism</i> , 35 , 178-184.
31	Sayed, M. D., Balbaa, S. I. and Afifi, M. S. (1973b). The lipid content of the seeds of <i>Citrullus colocynthis</i> . <i>Planta Medica</i> , 24 , 41-45.
32	Abudayeh Z. H. M., Lamazian H. R., Sereda P., Chekman I., Al Khalifa I. I., Al Azzam K. M. and Hassounah L. K. M. (2016). Comparative study of amino acid composition in the seeds, pulp and rind from <i>Citrullus colocynthis</i> fruits. <i>International Journal of Pharmacognosy and Phytochemical Research</i> , 8 (3), 433-437.
33	Kalhor, M. A., Afza, N., Saleem, M. and Malik, A. (2002). Pharmacochemical studies of the oil, aerial parts, pulp and peel of <i>Citrullus colocynthis</i> . <i>Journal of Chemical Society of Pakistan</i> , 24 , 274-276.
34	Gurudeeban, S., Ramanathan, T. and K. Satyavani. (2011). Characterization of volatile compounds from bitter apple (<i>Citrullus colocynthis</i>) using GC-MS. <i>International Journal of Chemical and Analytical Science</i> , 2 (8), 108-110.
35	Conservation and sustainable use of medicinal plants in Egypt, National Surveys (2016). UNDP, GEF, ASRT and NRC, Volumes 1-5.
36	Bailey, C. and Danin, A. (1981). Bedouin plant utilization in Sinai and the Negev. <i>Economic Botany</i> , 35 (2), 145-162.
37	Mahmoud, T. and Gairola, S. (2013). Traditional knowledge and use of medicinal plants in the Eastern Desert of Egypt: a case study from Wadi El-Gemal National Park. <i>Journal of Medicinal Plants Studies</i> , 1 (6), 10-17.
38	PDR for Herbal Medicines (1998). Medical Economic Co. Montvale, New Jersey, 639. ISBN 1563633612, 9781563633614.
39	https://www.webmd.com/vitamins/ai/ingredientmono-921/jimson-weed
40	Reza, H. J., Davoudi, A., Davoudi, F., Valizadegan, G., Goodarzi, H., Mahmoodi, S., Reza. M. G. and Faraji, M. (2013). <i>Citrullus colocynthis</i> as the cause of acute rectorrhagia. <i>Case Reports in Emergency Medicine</i> , Article ID 652192, 5. doi: 10.1155/2013/652192.
41	Qazan, W. S. H., Almasad, M. M. and Daradka, H. (2007). Short and long effects of <i>Citrullus colocynthis</i> L. on reproductive system and fertility in female Sprague Dawley rats. <i>Pak. J. Biol. Sci.</i> , 10 (16), 2699-2703.
42	Barghamdi, B., Ghorat, F., Asadollahi, K., Sayehmiri, K., Peyghambari, R. and Abangah, G. (2016). Therapeutic effects of <i>Citrullus colocynthis</i> fruit in patients with type II diabetes: A clinical trial study. <i>J. Pharm. Bioallied Sci.</i> , 8 (2), 130-134.
43	Jouad, H., Haloui, M., Rhiauani, H., El Hilaly, J. and Eddouks, M. (2001). Ethnobotanical survey of medicinal plants used for the treatment of diabetes, cardiac and renal diseases in the North centre region of Morocco (Fez-Boulemane). <i>Journal of Ethnopharmacology</i> , 2-3 (77), 175-182.
44	Khan, S. A., Shelleh, H. H., Bhat, A. R. and Bhat, K. S. (2003). Colocynth toxicity. A possible cause of bloody diarrhea. <i>Saudi Medical Journal</i> , 8 (24), 904-906.

45	Shafaei, A., Esmaili, H. Rad, S., Delazar, A. and Behjati, M. (2012). <i>Citrullus colocynthis</i> as a medicinal or poisonous plant: a revised fact. <i>JMPR.</i> , 35 (6), 4922–4927.
46	Mitchell, J. and Rook, A. (1979). Botanical Dermatology: Plants and Plant Products Injurious to the Skin. Greengrass, Vancouver, Canada. ISBN:0889780471, 9780889780477.
47	Blacow, N., Wade, W. and Red, A. (1972). Martindale: The Extra Pharmacopoeia. 26 th ed. London: <i>The Pharmaceutical Press</i> , 1627-1628.
48	Shah, A. H., Qureshi, S., Tariq, M. and Ageel, A. M. (1989). Toxicity studies on six plants used in the traditional Arab system of medicine. <i>Phytother. Res.</i> , 3 , 25-29.
49	Harnmarsten, G. and Lindgren, G. (1941-43). Ein Fall von Koloquinten-Vergiftung. <i>Vergiftungsfälle</i> 12 , A919: 107-110.
50	Pashmforosh, M., Vardanjani, H. R., Vardanjani, H. R., Pashmforosh, M. and Khodayar, M. J. (2018). Topical anti-inflammatory and analgesic activities of <i>Citrullus colocynthis</i> extract cream in rats. <i>Medicina</i> , 54 , 51- 61.
51	Marzouk, B., Marzouk, Z., Haloui, E., Fenina, N., Bouraoui, A. and Aouni, M. (2010). Screening of analgesic and anti-inflammatory activities of <i>Citrullus colocynthis</i> from southern Tunisia. <i>Journal of Ethnopharmacology</i> , 128 (1), 15-19.
52	Marzouk, B., Marzouk, Z., Fenina, N., Bouraoui, A. and Aouni, M. (2011). Anti-inflammatory and analgesic activities of Tunisian <i>Citrullus colocynthis</i> Schrad. Immature fruit and seed organic extracts. <i>Eur. Rev. Med. Pharmacol. Sci.</i> , 15 (6), 665-672.
53	Sanadgol, N., Najafi, S., Ghasemi, L. V., Motalleb, G and Estakhr, J. (2011). A study of the inhibitory effects of <i>Citrullus colocynthis</i> (CCT) using hydro-alcoholic extract on the expression of cytokines: TNF-and IL-6 in high fat diet-fed mice towards a cure for diabetes mellitus. <i>J. Pharmacognosy & Phytotherapy</i> , 3 (6), 81-88.
54	Akhzari, M., Mirghiasi, S., Vassaf, M., Bidgoli, M. and Tari, Z. (2015). The effect of <i>Citrullus colocynthis</i> on the reduction of inflammatory agents in osteoarthritis. <i>Mol. Biol.</i> , 4 (4), 33- 38.
55	Aly, A. M. and Naddaf, A. (2006). Anti-inflammatory activities of Colocynth topical gel. <i>J. Med. Sci.</i> , 6 , 216-221.
56	Rajamanickam, E., Gurudeeban, S., Ramanathan, T. and Satyavani, K. (2010). Evaluation of anti-inflammatory activity of <i>Citrullus colocynthis</i> . <i>Int. J. Cur. Res.</i> , 2 , 67-69.
57	Soufane, S., Bedda, A., Mahdeb, N. and Bouzidi, A. (2013). Acute toxicity study on <i>Citrullus colocynthis</i> fruit methanol extract in albino rats. <i>Journal of Applied Pharmaceutical Science</i> , 3 (6), 88-93.
58	Dehghani, F. and Panjehshahin, M. R. (2006). The toxic effect of alcoholic extract of <i>Citrullus colocynthis</i> on rat liver. <i>Iranian Journal of Pharmacology & Therapeutics</i> , 5 , 117-119.
59	Shaikh, J., Shaikh, D., Bin Rahman, A. and Shafi, S. (2016). Antimicrobial and toxicological studies on fruit pulp of <i>Citrullus colocynthis</i> L. <i>Pak. J. Pharm. Sci.</i> , 29 (1), 9-15.
60	Elalfy, M. M., Farag, A., Helmy, A. A., Metwaly, Z. E. and Ali, F. R. (2019). Hematological, biochemical and cytotoxic effect of ethanolic raw extract of Egyptian <i>Citrullus colocynthis</i> in sprague dawley rats. <i>Enz. Eng.</i> , 8 , 165.

61	Goldfain, D., Laverigne, A., Galian, A., Chauveinc, L. and Prudhomme, F. (1989). Peculiar acute toxic colitis after ingestion of colocynth: a clinicopathological study of three cases. <i>Gut</i> , 30 (10), 1412-1418.
62	Banerjee, S. P. and Dandiya, P. C. (1967). Smooth muscle and cardiovascular pharmacology of α -elaterin-2-D-glucopyranoside glycoside of <i>Citrullus colocynthis</i> . <i>J. Pharm. Sci.</i> , 56:1665-1667.
63	Lewin, L. (1962). Gifte und Vergiftungen. Lehrbuch der Toxikologie. Fiinfte unveranderte Ausgabe. UImIDonau: Karl F. Haug Verlag, 722-723.
64	Vohora, S. B., Khan, M. S. Y. and Afaq, S. H. (1973). Antifertility studies on Unani herbs. Part 2. Antioviulatory effects of 'hanzal', 'halun', 'kalonji' and 'sambhalu'. <i>Indian J. Pharm.</i> , 35 , 100-102.
65	Prakash, A. O., Saxena, V., Shukla, S., Tewari, R. K., Mathur, S., Gupta, A., Sharma, S. and Mathur, R. (1985). Anti-implantation activity of some indigenous plants in rats. <i>Acta Eur. Fertil.</i> , 16 (6), 441-448.
66	Osol, A. and Farrar, G. E. (1955). The Dispensatory of the United States of America. 25 th ed. Philadelphia: J.B. Lippincott Company, 359-360.
67	Anonymous. (1941). The Extra Pharmacopoeia - Martindale. 22 nd ed. London: The Pharmaceutical Press.
68	Sollmann, T. (1957). A Manual of pharmacology and its applications to therapeutics and toxicology. 8 th ed. Philadelphia: W.B. Saunders Company, 216.
69	Sweetman, S. C. (Ed). (2007). Martindale: The Complete Drug Reference. London: Pharmaceutical Press. Electronic version, London.
70	Agarwal, V., Sharma, A. K., Upadhyay, A., Singh, G. and Gupta, R. (2012). Hypoglycemic effects of <i>Citrullus colocynthis</i> roots. <i>Acta Pol Pharm.</i> , 69 (1), 75-79.
71	Huseini, H. F., Darvishzadeh, F., Heshmat, R., Jafariazar, Z., Raza, M. and Larijani, B. (2009). The clinical investigation of <i>Citrullus colocynthis</i> (L.) schrad fruit in treatment of Type II diabetic patients: a randomized, double blind, placebo controlled clinical trial. <i>Phytother. Res.</i> , 23 (8), 1186-1189.
72	Benariba, N., Djaziri, R., Zerriouh, B. H., Boucheri1, K., Louchami, K., Sener, A. and Willy, J. M. (2009). Antihyperglycemic effect of <i>Citrullus colocynthis</i> seed aqueous extracts in streptozotocin-induced diabetic rats. <i>Metabolic and Functional Research on Diabetes</i> , 2 , 71-77.
73	Jayaraman, P. N. N., Arihara, S., Anitha, T. and Joshi, V. D. (2009). Antidiabetic effect of petroleum ether extract of <i>Citrullus colocynthis</i> (L.) Schrad. fruits against streptozotocin-induced hyperglycemic rats. <i>Rom. J. Biol.-Plant Biol.</i> , 54 , 127-134.
74	Nmila, R., Gross, R., Rchid, H., Roye, M., Manteghetti, M., Petit, P., Tijane, M., Ribes, G. and Sauvaire, Y. (2000). Insulinotropic effect of <i>Citrullus colocynthis</i> fruit extracts. <i>Planta Medica</i> , 66 (5), 418-423.
75	Esmaeel, E., Somaieh, B., Mahmoud, H., Ghorban, M. Z., Saeed, S., Javad, J. (2016). Effect of hydroalcoholic leaves extract of <i>Citrullus colocynthis</i> on induction of insulin secretion from isolated rat islets of Langerhans. <i>Asian Pacific Journal of Tropical Disease</i> , 8 (6), 638-641.

76	Atole, S., Jangde, C., Philip, P., Rekhe, D., Aghav, D., Waghode, H. J. and Chougule, A. M. (2009). Safety evaluation studies of <i>Citrullus colocynthis</i> for diabetes in rats. <i>Veterinary World</i> , 2 (11), 423-425..
77	Nikbakht, M. and Gheatasi, I. (2006). Evaluation of the effect of hydroalcoholic extract of <i>Citrullus colocynthis</i> in normoglycemic and streptozocine (STZ) induced diabetic male rats. <i>Armaghane Danesh Bimonthly Journal</i> , 11 (2), 63-71.
78	Huseini, H. F., Zaree, A., Heshmat, R., Larijani, B., Fakhrzadeh, H., Rezaii Sharifabadi, R., Naderi, G. A., Zaringhalam, J. and Shikh Samani, A. H. (2006). The effect of <i>Citrullus colocynthis</i> (L.) Schrad. Fruit on oxidative stress parameters in type II diabetic patients. <i>Journal of Medicinal Plants</i> , 1 (17), 55-60.
79	Karimabad, M. N., Niknia, S., Golnabadi, M. B., Poor, S. F., Hajizadeh, M. R. and Mahmoodi, M. (2020). Effect of <i>Citrullus colocynthis</i> extract on glycated hemoglobin formation (<i>in vitro</i>). <i>The Eurasian Journal of Medicine</i> , 52 (1), 47-51.
80	Benariba, N., Djaziri, R., Zerriouh, B. H., Bellakhdar, W., Hupkens, E., Boucherit, Z. and Malaisse, W. J. (2012). Short-and long-term effects of various <i>Citrullus colocynthis</i> seed extracts in normal and streptozotocin-induced diabetic rats. <i>International Journal of Molecular Medicine</i> ; 30 (6), 1528-36.
81	Amin, A., Tahir, M. and Lone, K. P. (2017). Effect of <i>Citrullus colocynthis</i> aqueous seed extract on beta cell regeneration and intra-islet vasculature in alloxan induced diabetic male albino rats. <i>JPMMA The Journal of the Pakistan Medical Association</i> , 67 (5), 715-721.
82	Li, Y., Zheng, M., Zhai, X., Huang, Y., Khalid, A., Malik, A., Shah, P., Karim, S., Azhar, S. and Hou X. (2015). Effect of - <i>Gymnema sylvestre</i> , <i>Citrullus colocynthis</i> and <i>Artemisia absinthium</i> on blood glucose and lipid profile in diabetic human. <i>Acta. Pol. Pharm.</i> , 72 , 981-985.
83	Pooladvand, V., Taghavi, S., Mahmoodi, M., Tavakolian, F. V. and Hosseini, Z. M. (2011). Histological alterations due to the consumption of different doses of <i>Citrullus colocynthis</i> fruit in normal and diabetic male rats. <i>J. Mazandaran Univ. Med. Sci.</i> , 21 (82), 63-71.
84	Amin, A. and Tahir, M. (2016). Alpha cells a 'therapeutic target': effect of <i>Citrullus colocynthis</i> on alpha cell count in healthy and alloxan induced diabetic male albino rats. <i>World Journal of Pharmaceutical Research</i> , 11 (5), 329-339.
85	Abdel-Hassan, A., Abdel-Barry, J. A. and Mohammed, T. S. (2000). The hypoglycaemic and antihyperglycaemic effect of <i>Citrullus colocynthis</i> fruit aqueous extract in normal and alloxandiabetic rabbits. <i>J. Ethnopharmacol</i> , 71 , 325-330.
86	Ostovar, M., Akbari, A., Anbardar, M. H., Iraj, A., Salmanpour, M., Hafez, G. S., Heydari, M. and Shams, M. (2020). Effects of <i>Citrullus colocynthis</i> L. in a rat model of diabetic neuropathy. <i>Journal of Integrative Medicine</i> , 18 (1), 59-67.
87	Bourhia, M., Messaoudi, M., Bakrim, H., Mothana, R. A., Sddiqui, N. A., Almarfadi, O. M., El Mzibri, M., Gmouh, S., Laglaoui, A. and Benbacer, L. (2020). <i>Citrullus colocynthis</i> (L.) Schrad: Chemical characterization, scavenging and cytotoxic activities. <i>Open Chemistry</i> , 18 , 986-994.
88	Jayaraman, R. and Christina, A. J. M. (2013) Evaluation of <i>Citrullus colocynthis</i> fruits on <i>in vitro</i> antioxidant activity and <i>in vivo</i> DEN/PB induced hepatotoxicity. <i>Int. J. Applied Res. in Nat. Pro.</i> , 6 (1), 1-9.

89	Dallak, M. (2011). <i>In vivo</i> , hypolipidemic and antioxidant effects of <i>Citrullus colocynthis</i> pulp extract in alloxan induced diabetic rats. <i>Afr. J. Biotech.</i> , 10 (48), 9898-9903.
90	Gill, N. S., Kaur, S., Arora, R. and Bail, M. (2011). Screening of antioxidant and antiulcer potential of <i>Citrullus colocynthis</i> methanolic seed extract. <i>Res. J. Phytochemistry</i> , 5 (2), 98- 106.
91	Roy, R. K., Thakur, M. and Dixit, V. K. (2007). Effect of <i>Citrullus colocynthis</i> on hair growth in albino rats. <i>Pharma Bio.</i> , 45 (10), 739-744.
92	Dhanotia, R., Chauhan, N. S., Saraf D. K. and Dixit, V. K. (2011). Effect of <i>Citrullus colocynthis</i> Schrad fruits on testosterone-induced alopecia. <i>Nat. Prod. Res.</i> , 25 (15), 1432-1443.
93	Gurudeeban, S., Ramanathan, T. and Satyavani, K. (2010). Antioxidant and radical scavenging activity of <i>Citrullus colocynthis</i> . <i>Inventi Rapid: Nutraceuticals</i> , 1 , 38.
94	Rodge, S. V. and Biradar, S. D. (2012). Preliminary phytochemical screening and antimicrobial activity of <i>Citrullus colocynthis</i> (Linn.) Shared. <i>Indian J. Plant Sci.</i> , 2 (1), 19- 23.
95	Eidi, S., Azadi, H. G., Rahbar, N. and Mehmannaavaz, H. R. (2015). Evaluation of antifungal activity of hydroalcoholic extracts of <i>Citrullus colocynthis</i> fruit. <i>Journal of Herbal Medicine</i> , 5 (1), 36-40.
96	Archana, M., Garima, S., Sumita, K., Meenakshi, S. and Kothari, S. L. (2013). Antimycobacterial activity of <i>Citrullus colocynthis</i> (L.) Schrad. against drug sensitive and drug resistant Mycobacterium tuberculosis and MOTT clinical isolates. <i>J. Ethnopharm.</i> , 149 (1), 195-200.
97	Al-hejjaj, M. Y., Alhurba, Y. A. and Mohamad, S. A. (2010). Study of alkaloid extract from <i>Citrullus colocynthis</i> fruit and its antimicrobial activity screening. <i>J. Basrah. Res. (Sciences)</i> , 36 (4), 42-47.
98	Memon, U., Brohi, A. H., Ahmed, S. W., Azhar, I. and Bano, H., (2003). Antibacterial screening of <i>Citrullus colocynthis</i> . <i>Pak. J. Pharm. Sci.</i> , 16 (1), 1-6.
99	Menaria, K., Swarnakar, G. and Kumawat, A. (2020). Anthelmintic effect of <i>Citrullus colocynthis</i> on the tegument of <i>Cotylophoron cotylophorum</i> by light microscope. <i>IJPSR</i> , 11 (8), 4029-4038.
100	Swarnakar, G. and Kumawat, A. (2014). <i>In vitro</i> anthelmintic effect of <i>Citrullus colocynthis</i> on tegument of amphistome <i>Orthocoelium scolicoelium</i> (Trematoda: Digenea). <i>Int. J. Curr. Microbiol. App. Sci.</i> , 3 (6), 571-582.
101	Ahmed, C. N., Hamad, K. K. and Qadir, F. A. (2019). <i>Haemonchus contortus</i> as a model in assessing activity of <i>Citrullus colocynthis</i> fruit extract to control benzimidazole-resistant parasitic nematodes. <i>ZJPAS</i> , 31 (5): 61-70.
102	Damor, R. and Swarnakar, G. (2018). In vitro anthelmintic effects of fruit extracts of <i>Citrullus colocynthis</i> on liver fluke <i>Fasciola gigantica</i> in buffaloes. <i>International Journal of Innovative Research and Revie.</i> , 6 (1), 1-11.
103	Talole, B. B., Baheti, D. G. and More, P. A. (2013). <i>In vitro</i> helmintholytic activity of leaves of <i>Citrullus colocynthis</i> . <i>International Journal of Research in Pharmacy and Chemistry</i> , 3 (2), 240-243.
104	Grossman, S., Dovrat, S., Gottlieb, H. E. and Bergman, M. (2007). Growth inhibitory activity of cucurbitacin glucosides isolated from <i>Citrullus colocynthis</i> on human breast cancer cells. <i>Biochem. Pharmacol.</i> , 73 (1), 56-67.

105	Shokrzadeh, M., Chabra, A., Naghshvar, F. and Ahmadi, A. H. (2013). The mitigating effect of <i>Citrullus colocynthis</i> (L.) fruit extract against genotoxicity induced by cyclophosphamide in mice bone marrow cells. <i>The Scientific World Journal</i> , Article ID 980480: 8.
106	Ayyad, S. E1. , Abdel-Lateff, A., Alarif, W. M., Patacchioli, F. R., Badria, F. A. and Ezmirly, S. T. (2011). <i>In vitro</i> and <i>in vivo</i> study of cucurbitacins-type triterpene glucoside from <i>Citrullus colocynthis</i> growing in Saudi Arabia against hepatocellular carcinoma. <i>Environmental toxicology and pharmacology</i> , 33 (2), 245-251.
107	Sari, M., Nemmiche, S., Benmehdi, H., Amrouche, A., Hamadi, A. and Sari, D. C. (2019). Hypolipidemic and antioxidant effects of <i>Citrullus colocynthis</i> seeds oil in high-fat diets induced obese rats. <i>Phytotherapie</i> , 17 (6), 310-320.
108	Yazit, S. M., Nemmiche, S., Amamou, F., Meziane, R. K. and Chabane-Sari, D. (2020). Anti-hyperlipidemic effect of fatty acids methyl esters (FAMES) of <i>Citrullus colocynthis</i> in high-fat diet induced obesity in rats. <i>Phytothérapie</i> , 18 , 131-139.
109	Vakiloddin, S., Fuloria, N., Fuloria, S., Dhanaraj, S. A., Balaji, K. and Karupiah, S. (2015). Evidence of hepatoprotective and antioxidant effect of <i>Citrullus colocynthis</i> fruits in paracetamol induced hepatotoxicity. <i>Pak. J. Pharm. Sci.</i> , 28 (3), 951-957.
110	Iqbal, A. D., Sharma, V., Saxena, R. C. and Bansal, S. K. (2011). Hepatoprotective activity of <i>Citrullus colocynthis</i> Linn. <i>Inventi Rapid: Ethnopharmacolog</i> , 2 (2): ISSN 0976-3805.
111	Iqbal, A. D., Saxena, R. C. and Bansal, S. K. (2012). Hepatoprotection: A Hallmark of <i>Citrullus colocynthis</i> L. against paracetamol induced hepatotoxicity in swiss albino rats. <i>American Journal of Plant Sciences</i> , 7 , 1022-1027.